

M UELLER Implants & Periodontics

Thomas C. Mueller, DMD

Diplomate of the American Board of Periodontology

PRE-OPERATIVE INTRAVENOUS (IV) SEDATION INSTRUCTIONS

1. To reduce the chance of nausea, **DO NOT** eat or drink anything (including water) for at least 6 hours prior to your appointment.
 - a. If your surgery is in the morning, do not eat or drink anything between bedtime and your scheduled appointment.
 - b. If your surgery is in the afternoon, a light liquid breakfast before 7:00 AM is encouraged.
 - c. Unless specified by your dentist, all medicines taken on a routine basis should be continued without interruption. Please swallow with a minimal amount of water.
2. A responsible adult, over 18 years of age, **MUST** accompany you to the clinic and is encouraged to remain in the clinic during the entire procedure. Following the sedation, this adult must escort the patient home. A responsible adult should remain with the patient for the next 24 hours.
3. A parent or legal guardian **MUST** accompany minors or persons under the age of 18 years old.
4. For intravenous sedation, patient should wear clothing that is not restrictive to the neck or arms. Patients should wear loose-fitting tops of which sleeves can be rolled to the shoulder. Patients are requested to wear short-sleeved tops and button down front tops.
5. Contact lenses **MUST** be removed prior to sedation. Nail polish should be removed.
6. Following the anesthetic, patient should refrain from driving a vehicle or engaging in any activity that requires alertness for the next 24 hours. The patient cannot drive home.

**FAILURE TO FOLLOW ANY OF THESE INSTRUCTIONS WILL RESULT IN
CANCELLATION OF THE SEDATION APPOINTMENT**



2601 NW Rolling Green Dr., Corvallis, OR 97330 • Tel 541 757-8330 • Fax 541 757-0238

127 NE Eads St., Newport, OR 97365 • Tel 541 757-8330 • Fax 541 757-0238

